

Reprinted without change of pagination from
The Journal of Chemical Physics, Vol. 39, No. 3, p. 853, August 1, 1963

2p.

N 6 4 10 4 4 3 4

CODE NONE
(NASA CR-52462; TR-32-408)

Technical Report No. 32-408

Two-Electron, One- and Two-Center
Integrals

12
Murray Geller Repr. from
v. 39, no. 3, Aug. 1, 1963 p. 853 refs
NASA CR 52462

This paper presents results of one phase of research carried out at the Jet Propulsion Laboratory, California Institute of Technology, under Contract No. NAS 7-100, sponsored by the National Aeronautics and Space Administration.

(NASA Contract NAS 7-100)

jpl

Consent
JET PROPULSION LABORATORY
CALIFORNIA INSTITUTE OF TECHNOLOGY
PASADENA, CALIFORNIA

August, 1963

Reprint

Two-Electron, One- and Two-Center Integrals

MURRAY GELLER

Jet Propulsion Laboratory, Pasadena, California

(Received 14 February 1963)

IN a recent note,¹ Prosser and Blanchard mentioned the use of the Fourier convolution theorem method for the evaluation of one-electron, two-center integrals. This method has been applied by the author to one-electron integrals involving nonintegral Slater orbitals² and to one-electron integrals involving solid spherical harmonic operators.³

The present note is concerned with the application of this method to the evaluation of two-electron, two-center integrals⁴ (see Fig. 1) of the type

$$I = \int f(\mathbf{r}_{a1}) g(\mathbf{r}_{b2}) h(\mathbf{r}_{12}) d\mathbf{r}_1 d\mathbf{r}_2. \quad (1)$$

The integral is recovered by

$$I = (2\pi)^{-3} \int \tilde{f}(\mathbf{k}) \tilde{g}(\mathbf{k}) \tilde{h}(\mathbf{k}) \exp(-i\mathbf{k} \cdot \mathbf{R}) d\mathbf{k}, \quad (2)$$

where the bar indicates the appropriate Fourier transform

$$\tilde{\phi}(\mathbf{k}) = \int \exp(i\mathbf{k} \cdot \mathbf{r}) \phi(\mathbf{r}) d\mathbf{r}. \quad (3)$$

In the corresponding one-center case (the limit as \mathbf{R} goes to zero), the centers a and b coalesce (so that the subscripts a and b can be dropped) giving rise to the integral

$$J = \int f(\mathbf{r}_1) g(\mathbf{r}_2) h(\mathbf{r}_{12}) d\mathbf{r}_1 d\mathbf{r}_2, \quad (4)$$

which is recovered by

$$J = (2\pi)^{-3} \int \tilde{f}(\mathbf{k}) \tilde{g}(\mathbf{k}) \tilde{h}(\mathbf{k}) d\mathbf{k}. \quad (5)$$

As an example of Eqs. (4) and (5), Pitzer and Hameka⁵ have discussed the one-center integral

$$J = (32\pi)^{-2} \int r_1^4 \cos^2\theta_1 r_2^4 \cos^2\theta_2 (r_{12}^{-3} - 3z_{12}^2 r_{12}^{-5}) \times \exp(-r_1 - r_2) \sin\theta_1 \sin\theta_2 dr_1 dr_2 d\theta_1 d\theta_2 d\phi_1 d\phi_2. \quad (6)$$

The transform⁶ of $f(\mathbf{r}_1) = r_1^2 \cos^2\theta_1 \exp(-r_1)$ is given by

$$\tilde{f}(\mathbf{k}) = 32\pi(1+k^2)^{-4} [1 - k^2 - 4k^2 P_2(\cos u)]. \quad (7)$$

Since the function $g(\mathbf{r}_2)$ is identical, its transform is also given by Eq. (7). For the transform of




FIG. 1. Coordinate system.

$$h(\mathbf{r}_{12}) = r_{12}^{-3} - 3z_{12}^2 r_{12}^{-5} = -2P_2(\cos\theta_{12})/r_{12}^3, \quad (8)$$

we have

$$\tilde{h}(\mathbf{k}) = 8\pi P_2(\cos u)/3. \quad (9)$$

After substituting Eqs. (7) and (9) into Eq. (5), and integrating over u and v (the angular components of the \mathbf{k} vector), we find

$$J = \frac{32\pi}{105} \left[11 \int_0^\infty \frac{k^6}{(1+k^2)^8} dk - 7 \int_0^\infty \frac{k^4}{(1+k^2)^8} dk \right]. \quad (10)$$

When these two simple integrals of Eq. (10) are evaluated, the result $J = -\frac{1}{1680}$ is obtained in agreement with the result quoted by Pitzer and Hameka.⁵

An interesting point to notice is that this method does not involve the introduction of delta-function terms. The method is further not limited to one-center nor to specific forms for the f , g , and h functions.

The author wishes to acknowledge the informative discussions with Dr. Howard B. Levine of North American Aviation Science Center as to the range of validity of the Fourier convolution theorem.

¹ F. P. Prosser and C. H. Blanchard, J. Chem. Phys. **36**, 1112 (1962).

² M. Geller, J. Chem. Phys. **36**, 2424 (1962).

³ M. Geller, "Two-Center Integrals over Solid Spherical Harmonics," J. Chem. Phys. (to be published).

⁴ E. Meeron, J. Chem. Phys. **28**, 630 (1958), has given a general proof of the n -fold convolution integral.

⁵ R. M. Pitzer and H. F. Hameka, J. Chem. Phys. **37**, 2725 (1962).

⁶ See Ref. 3 for the method of evaluation of the transforms in detail.